

Lipoprotein(a) Reduction in Persons with Cardiovascular Disease

Sotirios Tsimikas, M.D., Ewa Karwatowska-Prokopczuk, M.D., Ph.D., Ioanna Gouni-Berthold, M.D., Jean-Claude Tardif, M.D., Seth J. Baum, M.D., Elizabeth Steinhagen-Thiessen, M.D., Michael D. Shapiro, D.O., Erik S. Stroes, M.D., Patrick M. Moriarty, M.D., Børge G. Nordestgaard, M.D., D.M.Sc., Shuting Xia, M.S., Jonathan Guerriero, M.B.A., Nicholas J. Viney, B.Sc., Louis O'Dea, M.B., B.Ch., B.A.O., and Joseph L. Witztum, M.D., for the AKCEA-APO(a)-L_{Rx} Study Investigators*

ABSTRACT

BACKGROUND

Lipoprotein(a) levels are genetically determined and, when elevated, are a risk factor for cardiovascular disease and aortic stenosis. There are no approved pharmacologic therapies to lower lipoprotein(a) levels.

METHODS

We conducted a randomized, double-blind, placebo-controlled, dose-ranging trial involving 286 patients with established cardiovascular disease and screening lipoprotein(a) levels of at least 60 mg per deciliter (150 nmol per liter). Patients received the hepatocyte-directed antisense oligonucleotide AKCEA-APO(a)-L_{Rx}, referred to here as APO(a)-L_{Rx} (20, 40, or 60 mg every 4 weeks; 20 mg every 2 weeks; or 20 mg every week), or saline placebo subcutaneously for 6 to 12 months. The lipoprotein(a) level was measured with an isoform-independent assay. The primary end point was the percent change in lipoprotein(a) level from baseline to month 6 of exposure (week 25 in the groups that received monthly doses and week 27 in the groups that received more frequent doses).

RESULTS

The median baseline lipoprotein(a) levels in the six groups ranged from 204.5 to 246.6 nmol per liter. Administration of APO(a)-L_{Rx} resulted in dose-dependent decreases in lipoprotein(a) levels, with mean percent decreases of 35% at a dose of 20 mg every 4 weeks, 56% at 40 mg every 4 weeks, 58% at 20 mg every 2 weeks, 72% at 60 mg every 4 weeks, and 80% at 20 mg every week, as compared with 6% with placebo (P values for the comparison with placebo ranged from 0.003 to <0.001). There were no significant differences between any APO(a)-L_{Rx} dose and placebo with respect to platelet counts, liver and renal measures, or influenza-like symptoms. The most common adverse events were injection-site reactions.

CONCLUSIONS

APO(a)-L_{Rx} reduced lipoprotein(a) levels in a dose-dependent manner in patients who had elevated lipoprotein(a) levels and established cardiovascular disease. (Funded by Akcea Therapeutics; ClinicalTrials.gov number, NCT03070782.)

The authors' affiliations are listed in the Appendix. Address reprint requests to Dr. Tsimikas at the Sulpizio Cardiovascular Center, Vascular Medicine Program, University of California, San Diego, 9500 Gilman Dr., La Jolla, California 92093, or at stsimikas@health.ucsd.edu.

*A list of the AKCEA-APO(a)-L_{Rx} Study Investigators is provided in the Supplementary Appendix, available at NEJM.org.

This article was published on January 1, 2020, at NEJM.org.

N Engl J Med 2020;382:244-55.

DOI: 10.1056/NEJMoa1905239

Copyright © 2020 Massachusetts Medical Society.

LIPOPROTEIN(A) IS COMPOSED OF A LOW-density lipoprotein (LDL)-like moiety bound covalently to apolipoprotein(a).^{1,2} Lipoprotein(a) potentially contributes to cardiovascular disease through proatherogenic effects of its LDL-like moiety, proinflammatory effects of its oxidized phospholipid content, and prothrombotic effects through its inactive, plasminogen-like protease domain on apolipoprotein(a). Mechanistic, epidemiologic, and genetic evidence reported over the past 20 years provides support for the idea that elevated plasma lipoprotein(a) is an independent genetic risk factor for cardiovascular disease and calcific aortic-valve stenosis.^{3,4} In contrast, genetically determined low levels of lipoprotein(a) (<30 mg per liter [<75 nmol per liter]) are associated with a decreased risk of cardiovascular disease but not of other non-cardiovascular disease adverse phenotypes.⁵

There are currently no approved pharmacologic therapies that specifically target lipoprotein(a). Antisense oligonucleotides (ASOs) inhibit the production of apolipoprotein(a) in the hepatocyte, the source of approximately 99% of plasma lipoprotein(a).⁶ Preclinical proof-of-concept studies have established that ASOs targeting hepatic *LPA* messenger RNA (mRNA) specifically reduce plasma levels of lipoprotein(a).^{7,8} Subsequent phase 1 and 2 studies of a non-hepatocyte-targeted, second-generation ASO showed lowering of lipoprotein(a) levels in healthy participants who had elevated lipoprotein(a), as well as in patients with established cardiovascular disease and elevated plasma levels of lipoprotein(a).^{9,10} Advances in directing ASOs to hepatocytes by conjugation with a triantennary *N*-acetylgalactosamine (GalNAc₃) moiety, a high-affinity ligand for the asialoglycoprotein receptor on the surface of hepatocytes, have resulted in large increases (by a factor of 15 to 30) in their potency,¹⁰ with implications for improvements in the side-effect profile and safety of ASOs.^{11,12} AKCEA-APO(a)-L_{Rx} — here referred to as APO(a)-L_{Rx} and previously called IONIS-APO(a)-L_{Rx} — is a GalNAc₃-conjugated 2'-methoxyethyl chimeric second-generation ASO drug targeted to *LPA* mRNA. In a phase 2a trial, APO(a)-L_{Rx} was shown to result in a dose-dependent reduction of 66 to 92% in circulating lipoprotein(a) in participants with elevated lipoprotein(a) levels.¹⁰ The long half-life of APO(a)-L_{Rx} (approximately 1 month) that was observed in that trial prompted us to con-

sider longer dosing intervals in the trial we report here.

METHODS

TRIAL DESIGN

We conducted this phase 2, dose-ranging, randomized, double-blind, placebo-controlled trial evaluating APO(a)-L_{Rx} at 30 sites in five countries. Patients were randomly assigned to one of five groups; within each group, randomization was performed in a 5:1 ratio (APO(a)-L_{Rx}:placebo) (Fig. S1 in the Supplementary Appendix, available with the full text of this article at NEJM.org). The patients in each group were given one of five regimens, with APO(a)-L_{Rx} or placebo administered subcutaneously, for a minimum of 6 months: APO(a)-L_{Rx} at a dose of 20 mg every 4 weeks, 40 mg every 4 weeks, 60 mg every 4 weeks, 20 mg every 2 weeks, or 20 mg every week or physiologic saline placebo. By week 25, the groups receiving monthly doses would have reached 6 months of exposure (after the week 21 dose); however, the groups receiving doses every 2 weeks or every week would have reached only 5.5 and 5.75 months of exposure, respectively. Therefore, the timing of the analysis of the primary efficacy end point was moved to week 27 for these groups.

To collect additional long-term safety and efficacy data, treatment was continued up to 1 year or until the last enrolled patient had reached 6 months of treatment. The post-treatment follow-up period lasted 16 weeks to account for the long half-life of the drug.¹¹ Site visits were scheduled to occur every 4 weeks to collect efficacy and safety data. Platelet counts and renal-function tests were performed every 2 weeks, and liver-function tests were performed every 2 weeks for the first 3 months and monthly thereafter throughout the treatment period. The platelet count was analyzed simultaneously by both central and local laboratories, and APO(a)-L_{Rx} or placebo could be administered only if recent (within 14 days) platelet-count results were available. Monitoring and stopping rules, which were prespecified in the protocol (available at NEJM.org), included threshold limits on platelet count and renal and liver function.

Akcea Therapeutics sponsored the trial and was responsible for its conduct and oversight, the collection and management of the data, and the statistical analyses and data interpretation.

The protocol was approved by the relevant health authorities, institutional review boards, and ethics committees. An academic author and an author who is an employee of the sponsor wrote the first draft and last submitted draft of the manuscript and vouch for the completeness and accuracy of the data and for fidelity of the trial to the protocol. All the authors participated in revising the manuscript.

ELIGIBILITY

Patients who were 18 to 80 years of age and had established cardiovascular disease and an elevated screening plasma lipoprotein(a) level (≥ 60 mg per deciliter [150 nmol per liter]) were eligible for enrollment. Patients who were being treated with lipid-lowering medications could be enrolled if they had been on a stable drug regimen for 4 weeks before screening and were expected to remain on that regimen during the trial. Exclusion criteria included acute coronary syndrome, major cardiac surgery, or stroke or transient ischemic attack within 6 months before screening; coronary, carotid, or peripheral revascularization, major noncardiac surgery, or lipoprotein apheresis within 3 months before screening; heart failure of New York Heart Association (NYHA) class IV; uncontrolled hypertension (systolic blood pressure, >160 mm Hg; or diastolic blood pressure, >100 mm Hg); an estimated glomerular filtration rate of less than 60 ml per minute; a ratio of urine protein (in milligrams) to creatinine (in grams) of 250 or greater; a ratio of urine albumin (in milligrams) to creatinine (in grams) of 100 or greater; an alanine aminotransferase or aspartate aminotransferase level exceeding twice the upper limit of the normal range; an alkaline phosphatase or total bilirubin level exceeding the upper limit of the normal range; a platelet count lower than the lower limit of the normal range; a history of major bleeding or high risk of bleeding diathesis; and use of anticoagulant drugs. Additional details, including those of the laboratory measurements, are provided in the Supplementary Appendix. Trial participants provided written informed consent.

END POINTS

The primary efficacy end point was the percent change in lipoprotein(a) level from baseline to the primary analysis time point at 6 months of exposure (week 25 or week 27) in each APO(a)-

L_{Rx} group as compared with the pooled placebo group. The unranked secondary end points, which were analyzed in the same way, were the percent change from baseline in LDL cholesterol level, the percent of patients with a plasma lipoprotein(a) level of 50 mg per deciliter (125 nmol per liter) or lower, the percent of patients with a plasma lipoprotein(a) level of 30 mg per deciliter (75 nmol per liter) or lower, the percent change from baseline in apolipoprotein B level, the percent change from baseline in the level of oxidized phospholipids on apolipoprotein B, and the percent change from baseline in the level of oxidized phospholipids on apolipoprotein(a).

STATISTICAL ANALYSIS

The power calculations are shown in the Supplementary Appendix. All efficacy analyses were performed with the full analysis set, defined as all patients who had undergone randomization and had received at least one dose of APO(a)- L_{Rx} or placebo. The primary end point and secondary end points for oxidized phospholipids on apolipoprotein B, oxidized phospholipids on apolipoprotein(a), LDL cholesterol, and apolipoprotein B were analyzed with the use of an analysis of covariance model with treatment groups as factors and the log-transformed baseline value for each respective measure as covariate. Responder analyses (i.e., in which each patient is considered as either having a response or having no response) were performed with the use of a logistic-regression model with the log-transformed baseline level of lipoprotein(a) as the concomitant variable. Missing data for the primary and secondary efficacy end points were handled with a multiple-imputation model containing baseline and postbaseline values, stratified according to treatment group. The imputations were performed for postbaseline values by the Markov chain Monte Carlo method. Supportive efficacy analyses over time, including primary and secondary efficacy data with data collected beyond the primary analysis time point, were performed with a mixed model of repeated measurements. Because of the exploratory nature of this phase 2 trial, the P values and widths of the 95% confidence intervals were not adjusted for multiplicity. All safety analyses were performed in the safety population, defined as all patients who underwent randomization and received at least one dose of APO(a)- L_{Rx} or placebo.

RESULTS

PATIENTS

Of the 459 patients who underwent assessment for eligibility, 286 were randomly assigned to one of five APO(a)-L_{Rx} regimens or placebo (Figs. S1 and S2). The first patient underwent randomization on March 27, 2017, and the last patient underwent randomization on January 16, 2018.

The mean (\pm SD) duration of treatment was 31.6 \pm 11.5 weeks (median, 32.1) among patients who received APO(a)-L_{Rx} and 31.2 \pm 12.0 weeks (median, 34.0) among those who received placebo. Approximately 60% of patients were younger than 65 years, and more than 30% were women (Table 1). By protocol design, all patients had established atherosclerotic cardiovascular disease, most commonly coronary artery disease; 4.5% had both carotid and peripheral artery disease. Between 35% and 54% of patients in each group had premature coronary artery disease, defined as a first cardiovascular event before 55 years of age for men and before 65 years of age for women. A majority of patients had a family history of coronary artery disease, and approximately one third had familial hypercholesterolemia. At trial entry, approximately 80 to 90% of the patients were receiving statin therapy, 50% were receiving ezetimibe, and 20% were receiving a PCSK9 inhibitor.

At baseline across all groups, median levels of lipoprotein(a) ranged from 205 to 247 nmol per liter (the upper limit of the normal range is 30 mg per deciliter [75 nmol per liter]),¹³ median levels of oxidized phospholipids on apolipoprotein B ranged from 20.3 to 24.6 nmol per liter (the top quartile in the general population is >8 nmol per liter), and median levels of oxidized phospholipids on apolipoprotein(a) ranged from 61.9 to 67.3 nmol per liter (the top quartile in the general population is >20 nmol per liter). Mean LDL cholesterol levels ranged from 67.6 to 89.3 mg per deciliter (1.75 to 2.31 mmol per liter). Mean estimated corrected LDL cholesterol levels ranged from 40.8 to 56.4 mg per deciliter, if it is assumed that 30% of lipoprotein(a) mass is cholesterol.¹⁴

MEAN PERCENT CHANGE IN LIPOPROTEIN(A) LEVEL

At 6 months of exposure (25 or 27 weeks), dose-dependent mean percent reductions in lipoprotein(a) from baseline were noted in all the

APO(a)-L_{Rx} groups, with decreases of 35% at a dose of 20 mg every 4 weeks, 56% at 40 mg every 4 weeks, 58% at 20 mg every 2 weeks, 72% at 60 mg every 4 weeks, and 80% at 20 mg every week, as compared with 6% for the pooled placebo group (P value range for the comparison with placebo, 0.003 to <0.001) (Fig. 1A). The two regimens of the same monthly dose (40 mg) — 40 mg every 4 weeks and 20 mg every 2 weeks — lowered lipoprotein(a) to similar extents. The lipoprotein(a)-lowering effect was noted within the first month and reached near-maximal effect by week 16 (Fig. 1B). In addition, dose-dependent absolute reductions in lipoprotein(a) levels were observed in all the APO(a)-L_{Rx} groups (Table 2). The lipoprotein(a) levels returned to baseline within 16 weeks after the last dose (Fig. S3), a finding consistent with previous observations.¹⁰ The mean percent changes in the lipoprotein(a) level in individual patients in each treatment group are shown in waterfall plots in Figure S4.

PERCENT OF PATIENTS ATTAINING PRESPECIFIED LIPOPROTEIN(A) LEVELS

The percent of patients with a lipoprotein(a) level of 50 mg per deciliter (125 nmol per liter) or lower at 6 months of exposure was 23% in the group that received 20 mg of APO(a)-L_{Rx} every 4 weeks, 63% in the group that received 40 mg every 4 weeks, 65% in the group that received 20 mg every 2 weeks, 81% in the group that received 60 mg every 4 weeks, and 98% in the group that received 20 mg every week. The corresponding odds ratios for this end point in comparison with the placebo group were 5.0 (95% confidence interval [CI], 1.2 to 21.0), 31.1 (95% CI, 7.3 to 131.4), 43.8 (95% CI, 9.8 to 195.0), 122.8 (95% CI, 24.0 to 627.4), and 1124.6 (95% CI, 109.3 to 11,571) (Fig. 1C).

The percent of patients with a lipoprotein(a) level of 30 mg per deciliter (75 nmol per liter) or lower at 6 months of exposure ranged from 6% in the group that received 20 mg every 4 weeks to 71% in the group that received 20 mg every week. The odds ratios for this end point in comparison with the placebo group were 7.3 (95% CI, 0.3 to 155.3) for 20 mg every 4 weeks, 27.9 (95% CI, 1.5 to 521.5) for 40 mg every 4 weeks, 59.9 (95% CI, 3.2 to 1128.0) for 20 mg every 2 weeks, 113.9 (95% CI, 6.2 to 2098.5) for 60 mg every 4 weeks, and 347.0 (95% CI, 18.3 to 6597.9) for 20 mg every week.

Table 1. Demographic, Clinical, and Laboratory Characteristics of the Patients at Baseline.*

Characteristic	APO(a)-Lrx				Pooled Placebo (N=47)
	20 mg Every 4 Wk (N=48)	40 mg Every 4 Wk (N=48)	20 mg Every 2 Wk (N=48)	60 mg Every 4 Wk (N=47)	
Age — yr	60.0±9.6	61.3±10.6	57.9±11.5	62.2±9.7	59.9±10.5
Age <65 yr — no. (%)	32 (67)	28 (58)	30 (62)	25 (53)	150 (63)
Age <50 yr — no. (%)	9 (19)	8 (17)	15 (31)	6 (13)	47 (20)
Female sex — no. (%)	19 (40)	12 (25)	17 (35)	14 (30)	82 (34)
White race — no./total no. (%)†	44 (92)	45 (94)	47 (98)	47 (100)	230 (96)
Cardiovascular risk factors — no. (%)					
Hypertension	28 (58)	35 (73)	34 (71)	31 (66)	153 (64)
Type 2 diabetes	7 (15)	8 (17)	5 (10)	4 (9)	10 (21)
Family history of coronary artery disease	34 (71)	31 (65)	33 (69)	29 (62)	161 (67)
Familial hypercholesterolemia	13 (27)	14 (29)	13 (27)	10 (21)	65 (27)
Current smoker	8 (17)	4 (8)	2 (4)	2 (4)	1 (2)
Body-mass index‡	29.0±4.51	27.2±3.52	28.7±4.18	28.5±3.96	28.4±4.25
Coronary artery disease — no. (%)	44 (92)	45 (94)	45 (94)	46 (98)	224 (94)
Premature coronary artery disease§	21 (44)	26 (54)	17 (35)	18 (38)	105 (44)
Myocardial infarction	25 (52)	25 (52)	31 (65)	20 (43)	128 (54)
Carotid artery disease — no. (%)	13 (27)	12 (25)	10 (21)	11 (23)	60 (25)
Peripheral artery disease — no. (%)	9 (19)	10 (21)	4 (8)	1 (2)	29 (12)
Stroke or TIA — no. (%)	6 (12)	6 (12)	7 (15)	4 (9)	28 (12)
Aortic-valve stenosis of grade 2 or higher — no. (%)	5 (10)	3 (6)	3 (6)	1 (2)	15 (6)
History of lipoprotein apheresis — no. (%)	1 (2)	2 (4)	2 (4)	1 (2)	8 (3)

Lipid-lowering therapy — no. (%)	42 (88)	44 (92)	43 (90)	44 (94)	44 (92)	217 (91)	39 (83)
Statin therapy							
High intensity	36 (75)	31 (65)	30 (62)	37 (79)	32 (67)	166 (69)	32 (68)
Moderate or low intensity¶	6 (12)	13 (27)	13 (27)	7 (15)	12 (25)	51 (21)	7 (15)
Ezetimibe	25 (52)	25 (52)	23 (48)	19 (40)	23 (48)	115 (48)	23 (49)
PCSK9 inhibitor	8 (17)	11 (23)	9 (19)	10 (21)	13 (27)	51 (21)	10 (21)
Platelet aggregation inhibitors — no. (%)	47 (98)	47 (98)	46 (96)	47 (100)	48 (100)	235 (98)	43 (91)
Median lipoprotein(a) level (IQR) — nmol/liter	246.6 (179.2–300.8)	220.0 (176.5–283.3)	238.2 (183.7–298.4)	204.5 (163.8–286.5)	233.7 (193.1–275.3)	224.3 (177.2–286.9)	231.6 (194.9–317.7)
Median OxPL-apoB level (IQR) — nmol/liter	24.6 (18.1–33.1)	23.1 (16.2–32.5)	23.9 (17.9–29.2)	20.3 (16.6–28.5)	23.7 (17.2–30.7)	23.3 (17.4–30.5)	21.2 (17.2–31.5)
Median OxPL-apo(a) level (IQR) — nmol/liter	66.3 (57.8–75.0)	65.9 (56.6–71.9)	67.3 (60.8–73.2)	61.9 (53.4–72.7)	67.1 (60.0–74.6)	65.8 (58.6–73.8)	69.2 (59.6–76.5)
Total cholesterol level — mg/dl	166.0±38.6	154.4±52.1	146.7±36.3	142.9±32.4	154.4±34.7	154.4±39.8	154.4±33.6
LDL cholesterol level — mg/dl	89.3±37.1	77.4±39.5	74.4±28.8	67.6±28.3	76.1±28.4	77.0±33.3	79.4±29.2
Estimated LDL-Ccorr — mg/dl	56.4±39.5	49.8±39.1	45.2±27.6	40.8±28.3	46.7±30.1	47.8±33.5	47.6±30.1
Apolipoprotein B level — mg/dl	80.7±23.6	71.9±23.4	69.3±19.8	68.5±18.8	70.6±19.2	72.2±21.3	73.8±16.9
HDL cholesterol level — mg/dl	54.1±15.5	54.1±19.3	54.1±19.3	50.3±11.6	58.0±19.3	54.1±19.3	50.3±19.3
Median triglyceride level (IQR) — mg/dl	97 (44–230)	97 (35–283)	106 (35–204)	106 (53–567)	89 (35–266)	97 (35–567)	106 (35–576)
hsCRP level — mg/liter	2.9±5.3	2.3±4.5	1.6±2.5	2.0±2.5	2.2±4.4	2.2±4.0	2.4±4.4

* Plus-minus values are means ±SD. To convert the values for lipoprotein(a) to nanomoles per liter, multiply by 2.5. To convert the values for cholesterol to millimoles per liter, multiply by 0.02586. To convert the values for triglycerides to millimoles per liter, multiply by 0.01129. HDL denotes high-density lipoprotein, hsCRP high-sensitivity C-reactive protein, IQR interquartile range, LDL low-density lipoprotein, OxPL-apoB oxidized phospholipids on apolipoprotein B, OxPL-apo(a) oxidized phospholipids on apolipoprotein(a), and TIA transient ischemic attack.

† Race was reported by the patient.

‡ Body-mass index is the weight in kilograms divided by the square of the height in meters.

§ Premature coronary artery disease was defined as disease that occurred before the age of 55 years in men and before the age of 65 years in women.

¶ This category includes patients who were not treated with statins.

|| The LDL cholesterol level corrected for lipoprotein(a) (LDL-Ccorr) was calculated as the LDL cholesterol level minus the lipoprotein(a) mass multiplied by 0.3.




Figure 1. Effect of APO(a)-L_{Rx} on Lipoprotein(a) Level.

Panel A shows the least-squares mean percent changes from baseline to the primary analysis time point (PAT) (i.e., 6 months of exposure [at week 25 in the groups that received monthly doses and at week 27 in the groups that received more frequent doses]). I bars denote 95% confidence intervals. Panel B shows the least-squares mean percent changes from baseline in lipoprotein(a) over time. Error bars denote 95% confidence intervals. Panel C shows the percent of patients with lipoprotein(a) levels of less than 50 mg per deciliter (125 nmol per liter) in each group at the primary analysis time point.

OTHER PRESPECIFIED MEASURES

At 6 months of exposure, we observed mean reductions in levels of oxidized phospholipids on apolipoprotein B and levels of oxidized phospholipids on apolipoprotein(a) in all APO(a)-L_{Rx} groups. The mean percent reductions in oxidized phospholipids on apolipoprotein B were 37% at a dose of 20 mg every 4 weeks, 57% at 40 mg every 4 weeks, 64% at 20 mg every 2 weeks, 79% at 60 mg every 4 weeks, and 88% at 20 mg every week, as compared with a 14% increase in the placebo group. The ratios of geometric mean reductions from baseline (APO(a)-L_{Rx}:placebo) were 0.55 (95% CI, 0.38 to 0.81), 0.37 (95% CI, 0.26 to 0.54), 0.32 (95% CI, 0.22 to 0.46), 0.18 (95% CI, 0.13 to 0.27), and 0.11 (95% CI, 0.07 to 0.16) in the groups that received 20 mg every 4 weeks, 40 mg every 4 weeks, 20 mg every 2 weeks, 60 mg every 4 weeks, and 20 mg every week, respectively. (A ratio of geometric means of <1 indicates a larger reduction from baseline in the APO(a)-L_{Rx} group than in the placebo group.)

The mean percent reductions in oxidized phospholipids on apolipoprotein(a) were 28% at a dose of 20 mg every 4 weeks, 49% at 40 mg every 4 weeks, 45% at 20 mg every 2 weeks, 63% at 60 mg every 4 weeks, and 70% at 20 mg every week, as compared with a 20% decrease in the placebo group. The ratios of the geometric mean reductions from baseline (APO(a)-L_{Rx}:placebo) were 0.91 (95% CI, 0.68 to 1.21), 0.64 (95% CI, 0.48 to 0.86), 0.69 (95% CI, 0.52 to 0.92), 0.46 (95% CI, 0.35 to 0.62), and 0.38 (95% CI, 0.28 to 0.51) for 20 mg every 4 weeks, 40 mg every 4 weeks, 20 mg every 2 weeks, 60 mg every 4 weeks, and 20 mg every week, respectively.

Table 2. Absolute Change from Baseline at 6 Months of Exposure.*

Measure	APO(a)-L _{Rx}					Pooled Placebo (N=47)
	20 mg Every 4 Wk (N=48)	40 mg Every 4 Wk (N=48)	20 mg Every 2 Wk (N=48)	60 mg Every 4 Wk (N=47)	20 mg Every Wk (N=48)	
Lipoprotein(a) — nmol/liter	-95.9±94.4	-116.9±71.7	-130.3±66.1	-149.5±67.4	-187.8±80.3	-15.2±34.6
Lipoprotein(a) — mg/dl	-38.4±7.7	-46.8±28.7	-52.1±26.4	-59.8±27.0	-75.1±32.1	-6.1±13.8
OxPL-apoB — nmol/liter	-8.0±10.3	-11.3±11.0	-12.2±7.9	-14.9±10.3	-20.1±8.5	3.7±8.1
OxPL-apo(a) — nmol/liter	-16.8±14.3	-24.5±20.1	-25.9±17.2	-33.3±16.8	-41.6±16.5	-12.3±14.7
LDL cholesterol — mg/dl	-5.6±27.4	-13.5±30.1	-13.2±19.8	-8.2±17.3	-16.4±14.8	-1.2±17.8
Apolipoprotein B — mg/dl	-2.2±17.4	-8.3±18.2	-6.3±11.6	-3.9±13.5	-10.9±10.9	0.6±12.0
Total cholesterol — mg/dl	-3.9±32.1	-11.6±32.1	-11.6±24.4	-3.9±23.2	-11.6±20.9	-3.9±21.3
HDL cholesterol — mg/dl	0.0±6.2	0.0±9.7	3.7±8.9	3.7±11.6	3.7±10.1	0.0±6.6
Triglycerides — mg/dl	-8.9±32.8	-8.9±31.0	0.0±52.3	0.0±50.5	-8.9±41.6	0.0±51.4
hsCRP — mg/liter	-0.9±4.24	-0.7±4.24	-0.3±2.84	-0.5±2.22	-0.1±6.30	-0.8±5.13

* Plus-minus values are means ±SD. The primary analysis time point was at 6 months of exposure: week 25 in the groups that received monthly doses and week 27 in the groups that received more frequent doses. To convert the values for cholesterol to millimoles per liter, multiply by 0.02586. To convert the values for triglycerides to millimoles per liter, multiply by 0.01129.

4 weeks, and 20 mg every week, respectively (Table 2 and Fig. S5A and S5B). We found mean reductions from baseline to 6 months in apolipoprotein B levels (ideal levels are <80 mg per deciliter in high-risk patients) and in laboratory-measured LDL cholesterol levels in all treatment groups (Table 2 and Fig. S5C and S5D). Waterfall plots of individual patients' percent changes from baseline in levels of oxidized phospholipids on apolipoprotein B, oxidized phospholipids on apolipoprotein(a), apolipoprotein B, and LDL cholesterol are shown in Figures S6 through S9.

SAFETY ANALYSES

Adverse events occurred in 90% of patients receiving APO(a)-L_{Rx} and in 83% of those receiving placebo during the treatment period; most of the events were mild or moderate (Table 3). Serious adverse events occurred in 10% of the patients receiving APO(a)-L_{Rx} and in 2% of those receiving placebo. Neither the overall incidence of adverse events nor the incidence of serious adverse events showed a dose-dependent pattern (Table 3). Overall, 5% of patients who received APO(a)-L_{Rx} and 4% of those who received placebo discontinued participation in the trial owing to adverse events. The most frequent adverse

events leading to discontinuation among patients who received APO(a)-L_{Rx} were myalgia or arthralgia or postinjection general discomfort (malaise). No patients discontinued participation for protocol-defined reasons. Injection-site reactions, the most frequently reported adverse event, occurred in 27% of the patients who received APO(a)-L_{Rx} and in 6% of those who received placebo during the trial. Approximately 7% of injections were associated with injection-site reactions irrespective of the dose or regimen (Table 3). These reactions were mostly mild, and the most common was erythema (26%). One patient discontinued treatment with APO(a)-L_{Rx} because of injection-site reactions.

Other adverse events that occurred in more than 10% of patients who received APO(a)-L_{Rx} and were more frequent than in the placebo group were urinary tract infection (13% vs. 6%), myalgia (12% vs. 11%), and headache (11% vs. 8%). The incidence of influenza-like symptoms was similar in the APO(a)-L_{Rx} group and the placebo group (7% and 6%, respectively). Platelet-count monitoring that was conducted every 2 weeks did not reveal a dose- or time-dependent effect of APO(a)-L_{Rx}, and no patient had a platelet count lower than 100,000 per cubic millimeter (Table 3

Table 3. Adverse Events and Laboratory Measurements during the Treatment Period.*

Event or Measurement	APO(a)-LRx				Pooled Placebo (N = 47)
	20 mg Every 4 Wk (N = 48)	40 mg Every 4 Wk (N = 48)	20 mg Every 2 Wk (N = 48)	20 mg Every 4 Wk (N = 47)	
Adverse events					
Any adverse event — no. of patients (%)	46 (96)	42 (88)	41 (85)	43 (91)	42 (88)
Mild	26 (54)	21 (44)	25 (52)	18 (38)	21 (44)
Moderate	14 (29)	18 (38)	14 (29)	20 (43)	19 (40)
Severe	6 (12)	3 (6)	2 (4)	5 (11)	2 (4)
Serious adverse event — no. of patients (%)	6 (12)	7 (15)	3 (6)	6 (13)	3 (6)
Adverse event leading to treatment discontinuation — no. of patients (%)	2 (4)	0	1 (2)	3 (6)	6 (12)
Death — no. of patients (%)	0	0	0	1 (2)	1 (2)
Injection-site reaction — no. of patients (%) †	5 (10)	15 (31)	11 (23)	12 (26)	22 (46)
Injections leading to injection-site reaction — no. of injections/total no. (%)	6/393 (1.5)	36/438 (8.2)	59/819 (7.2)	34/396 (8.6)	126/1495 (8.4)
Influenza-like symptoms — no. of patients (%) ‡	0	4 (8)	4 (8)	6 (13)	2 (4)
Laboratory measurements — no. of patients (%) §					
Platelet count					
<140,000/mm ³ ¶	3 (6)	8 (17)	3 (6)	3 (6)	8 (17)
<100,000/mm ³	0	0	0	0	0
ALT level >3x ULN	0	0	0	0	0
AST level >3x ULN	0	0	1 (2)	0	0
eGFR decrease of >25% from baseline	2 (4)	1 (2)	0	1 (2)	3 (6)
eGFR decrease of >40% from baseline	1 (2)	0	0	0	0
Urine albumin (mg):creatinine (g) ratio >250	1 (2)	0	0	1 (2)	1 (2)
Urine protein (mg):creatinine (g) ratio >500	1 (2)	0	0	2 (4)	1 (2)
24-hr creatinine clearance decrease of >40% from baseline**	0	1 (2)	1 (2)	0	0
24-hr creatinine clearance <45 ml/min/1.73 m ²	0	0	0	0	0
24-hr urine protein >1 g	0	0	0	0	0

* The treatment period was defined as the period from the first dose through one dosing interval after the last dose. ALT denotes alanine aminotransferase, AST aspartate aminotransferase, and ULN upper limit of the normal range.
 † Category includes any adverse event at the injection site persisting for at least 2 days.
 ‡ Category includes any of the following adverse events starting on the day of the injection or the next day: influenza-like illness, chills, myalgia, arthralgia, pyrexia, feeling hot, or a body temperature increase.
 § Values were confirmed by a second measurement within 1 week. In cases in which a second measurement was not available, the result was considered confirmed.
 ¶ Category includes patients who had two occurrences of the platelet count.
 || The estimated glomerular filtration rate (eGFR) was calculated with the Chronic Kidney Disease Epidemiology Collaboration formula.
 ** This event was due to prerenal causes in the patient who received 40 mg every 4 weeks and was due to incomplete 24-hr urine collection in the patient who received 20 mg every 2 weeks.

and Fig. S10). No patient had liver or renal toxic effects that met the predetermined thresholds for treatment discontinuation. There were no clinically significant changes in other laboratory measures (including coagulation panel), vital signs, or electrocardiographic measures. High-sensitivity C-reactive protein (CRP) levels were measured as a safety outcome to gauge potential proinflammatory effects of APO(a)-L_{Rx}. Baseline high-sensitivity CRP levels were between 2 and 3 mg per liter across the groups (Table 1). The absolute mean changes in the high-sensitivity CRP level at 6 months of exposure were -0.9 ± 4.2 mg per liter in the group that received 20 mg every 4 weeks, -0.7 ± 4.2 mg per liter in the group that received 40 mg every 4 weeks, -0.3 ± 2.8 mg per liter in the group that received 20 mg every 2 weeks, -0.5 ± 2.2 mg per liter in the group that received 60 mg every 4 weeks, and -0.1 ± 6.3 mg per liter in the group that received 20 mg every week, as compared with -0.8 ± 5.2 mg per liter in the pooled placebo group.

Two deaths occurred during the trial, both in patients receiving APO(a)-L_{Rx}. One patient in the group that received 60 mg every 4 weeks died in a road traffic accident, and one patient in the group that received 20 mg every week committed suicide as a result of depression.

DISCUSSION

APO(a)-L_{Rx} treatment resulted in dose-dependent reductions in lipoprotein(a) levels in patients with cardiovascular disease; these reductions were significant at all doses studied, with a mean 80% reduction at the highest dose (20 mg weekly). At the highest cumulative dose regimen, which was equivalent to 80 mg monthly, 98% of patients attained a lipoprotein(a) level of 50 mg per deciliter (125 nmol per liter) or lower, a target value supported by European¹⁵ and U.S.¹⁶ guidelines and by empirical data from patients treated with statins.¹⁷ We found reductions in levels of oxidized phospholipids on apolipoprotein B and oxidized phospholipids on apolipoprotein(a), both of which are proinflammatory components that are present on lipoprotein(a) and on apolipoprotein(a) and are linked to a higher atherothrombotic risk.¹⁸⁻²⁰ Finally, we noted reductions in LDL cholesterol and apolipoprotein B in patients receiving APO(a)-L_{Rx}, beyond those achieved with aggressive background lipid-lowering therapy.

The patients enrolled in this trial all had established cardiovascular disease, even though two thirds of the patients were under 65 years of age, which is generally younger than is typical in cardiovascular trials. This age profile was consistent with the lipoprotein(a) level being genetically determined and therefore a lifelong risk factor. Other ways in which the patients in the pooled trial population differed from participants in other cardiovascular trials were the relatively high percentages of patients with premature cardiovascular disease (40%) and familial hypercholesterolemia (approximately 30%) and the relatively low body-mass index and likelihood of having type 2 diabetes. That 97% of the patients were white is a limitation of the trial.

Patients with elevated lipoprotein(a) levels often cannot reach very low LDL cholesterol levels even with aggressive LDL-lowering therapy, because lipoprotein(a) cholesterol is comeasured with LDL cholesterol.²¹ Consistent with this observation is the fact that, in the current trial, mean LDL cholesterol levels at baseline were approximately 70 to 80 mg per deciliter despite treatment with up to three drugs. In the Further Cardiovascular Outcomes Research with PCSK9 Inhibition in Subjects with Elevated Risk (FOURIER) trial, patients in whom very low LDL cholesterol levels could not be achieved had markedly elevated lipoprotein(a) levels.²² Most LDL-lowering drugs do not substantially lower lipoprotein(a), and statins often have a neutral or modest lipoprotein(a)-increasing effect.^{17,23}

Evidence from studies of primary prevention indicate that elevated lipoprotein(a) levels are associated with an increased atherothrombotic risk — in particular, an increased risk of myocardial infarction.⁴ The role of lipoprotein(a) in the context of secondary prevention remains controversial. Although meta-analyses have intrinsic limitations, we would note that in a patient-level meta-analysis of statin outcomes trials that included 5751 events and 95,576 person-years at risk, the risk of elevated lipoprotein(a) was almost linearly associated with the risk of cardiovascular disease, and lipoprotein(a) levels greater than 50 mg per deciliter (125 nmol per liter) among patients who were taking statins were associated with significantly higher risk.¹⁷ Furthermore, a recent analysis of data from the placebo group in the FOURIER trial showed that higher levels of lipoprotein(a) are associated with

an increased risk of cardiovascular events in patients with established cardiovascular disease, irrespective of LDL cholesterol levels.²⁴

Among lipoproteins, lipoprotein(a) is the highest-capacity carrier of oxidized phospholipids,²⁵ which are responsible for many of its proinflammatory effects.²⁶⁻²⁸ In our trial, dose-dependent reductions were noted in oxidized phospholipids on apolipoprotein B and oxidized phospholipids on apolipoprotein(a) in plasma which, we posit, reflects effects of APO(a)-L_{Rx} on the oxidized phospholipid content of lipoprotein(a). In previous studies, oxidized phospholipids on apolipoprotein B, primarily reflecting the oxidized phospholipid content of lipoprotein(a), predicted first and recurrent myocardial infarction, stroke, and aortic stenosis.¹⁸⁻²⁰

The reduction in LDL cholesterol and apolipoprotein B levels by ASOs targeting *LPA* mRNA, in the presence of aggressive lipid-lowering therapy, has been documented in previous studies.^{9,10} Such findings may indicate that, when the synthesis of apolipoprotein(a) is inhibited in the liver, apolipoprotein B lipoproteins — which would otherwise be destined to become lipo-

protein(a), with relatively low plasma clearance as a result of weak interactions with the LDL receptor — are converted to LDL particles with relatively strong affinity for the LDL receptor and hence a relatively quick and efficient clearance from the blood.¹⁰

We did not observe marked changes in platelet, renal, or liver function, nor a between-group difference in the risk of influenza-like symptoms. The most common adverse events among patients who received APO(a)-L_{Rx} were injection-site reactions, which were generally mild.

Elevated levels of lipoprotein(a) are a cardiovascular risk factor for which no effective pharmacological therapy currently exists. In this trial, we found that APO(a)-L_{Rx} provided potent reductions in levels of lipoprotein(a) in patients with cardiovascular disease.

A data sharing statement provided by the authors is available with the full text of this article at NEJM.org.

Supported by Akcea Therapeutics.

Disclosure forms provided by the authors are available with the full text of this article at NEJM.org.

We thank the following employees of Akcea Therapeutics: Chelsey Jensen and Philip Piscitelli for overseeing the conduct of the trial, Eunju Hurh for critical data analysis, and Victoria Bartlett for project management.

APPENDIX

The authors' affiliations are as follows: the Divisions of Cardiovascular Medicine (S.T.) and Endocrinology and Metabolism (J.L.W.), University of California, San Diego, La Jolla, and Ionis Pharmaceuticals, Carlsbad (S.T., S.X., N.J.V.) — both in California; Akcea Therapeutics, Boston (E.K.-P., J.G., L.O.); Polyclinic for Endocrinology, Diabetes and Preventive Medicine, University of Cologne, Cologne, Germany (I.G.-B.); Montreal Heart Institute, Université de Montréal, Montreal (J.-C.T.); Excel Medical Clinical Trials, Boca Raton, FL (S.J.B.); the Department of Endocrinology and Metabolism, Charité-Universitätsmedizin Berlin, Freie Universität Berlin, Humboldt-Universität Berlin, Berlin Institute of Health, Berlin (E.S.-T.), and the Division of Geriatrics, University Medicine Greifswald, Greifswald (E.S.-T.) — both in Germany; the Center for Preventive Cardiology, Knight Cardiovascular Institute, Oregon Health and Science University, Portland (M.D.S.); the Department of Vascular Medicine, Academic Medical Center, Amsterdam (E.S.S.); the Division of Clinical Pharmacology, Department of Internal Medicine, University of Kansas Medical Center, Kansas City (P.M.M.); and the Department of Clinical Biochemistry and the Copenhagen General Population Study, Herlev and Gentofte Hospital, Copenhagen University Hospital, Herlev (B.G.N.), and the Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen (B.G.N.) — all in Denmark.

REFERENCES

- Berg K. A new serum type system in man — the Lp system. *Acta Pathol Microbiol Scand* 1963;59:369-82.
- Schmidt K, Noureen A, Kronenberg F, Utermann G. Structure, function, and genetics of lipoprotein (a). *J Lipid Res* 2016; 57:1339-59.
- Tsimikas S, Fazio S, Ferdinand KC, et al. NHLBI working group recommendations to reduce lipoprotein(a)-mediated risk of cardiovascular disease and aortic stenosis. *J Am Coll Cardiol* 2018;71:177-92.
- Nordestgaard BG, Langsted A. Lipoprotein (a) as a cause of cardiovascular disease: insights from epidemiology, genetics, and biology. *J Lipid Res* 2016;57: 1953-75.
- Emdin CA, Khera AV, Natarajan P, et al. Phenotypic characterization of genetically lowered human lipoprotein(a) levels. *J Am Coll Cardiol* 2016;68:2761-72.
- Crooke ST, Witztum JL, Bennett CF, Baker BF. RNA-targeted therapeutics. *Cell Metab* 2018;27:714-39.
- Merki E, Graham MJ, Mullick AE, et al. Antisense oligonucleotide directed to human apolipoprotein B-100 reduces lipoprotein(a) levels and oxidized phospholipids on human apolipoprotein B-100 particles in lipoprotein(a) transgenic mice. *Circulation* 2008;118:743-53.
- Merki E, Graham M, Taleb A, et al. Antisense oligonucleotide lowers plasma levels of apolipoprotein (a) and lipoprotein (a) in transgenic mice. *J Am Coll Cardiol* 2011;57:1611-21.
- Tsimikas S, Viney NJ, Hughes SG, et al. Antisense therapy targeting apolipoprotein(a): a randomised, double-blind, placebo-controlled phase 1 study. *Lancet* 2015;386:1472-83.
- Viney NJ, van Capelleveen JC, Geary RS, et al. Antisense oligonucleotides targeting apolipoprotein(a) in people with raised lipoprotein(a): two randomised, double-blind, placebo-controlled, dose-ranging trials. *Lancet* 2016;388:2239-53.

11. Yu RZ, Gunawan R, Post N, et al. Disposition and pharmacokinetics of a GalNAc₃-conjugated antisense oligonucleotide targeting human lipoprotein (a) in monkeys. *Nucleic Acid Ther* 2016;26:372-80.
12. Crooke ST, Baker BF, Xia S, et al. Integrated assessment of the clinical performance of GalNAc₃-conjugated 2'-O-methoxyethyl chimeric antisense oligonucleotides. I. human volunteer experience. *Nucleic Acid Ther* 2019;29:16-32.
13. Kostner GM, Avogaro P, Cazzolato G, Marth E, Bittolo-Bon G, Qunici GB. Lipoprotein Lp(a) and the risk for myocardial infarction. *Atherosclerosis* 1981;38:51-61.
14. Viney NJ, Yeang C, Yang X, Xia S, Witztum JL, Tsimikas S. Relationship between "LDL-C", estimated true LDL-C, apolipoprotein B-100, and PCSK9 levels following lipoprotein(a) lowering with an antisense oligonucleotide. *J Clin Lipidol* 2018;12:702-10.
15. Catapano AL, Graham I, De Backer G, et al. 2016 ESC/EAS guidelines for the management of dyslipidaemias. *Eur Heart J* 2016;37:2999-3058.
16. Grundy SM, Stone NJ, Bailey AL, et al. 2018 AHA/ACC/AACVPR/AAPA/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA guideline on the management of blood cholesterol: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol* 2019;73:3168-209.
17. Willeit P, Ridker PM, Nestel PJ, et al. Baseline and on-statin treatment lipoprotein(a) levels for prediction of cardiovascular events: individual patient-data meta-analysis of statin outcome trials. *Lancet* 2018;392:1311-20.
18. Tsimikas S, Willeit P, Willeit J, et al. Oxidation-specific biomarkers, prospective 15-year cardiovascular and stroke outcomes, and net reclassification of cardiovascular events. *J Am Coll Cardiol* 2012;60:2218-29.
19. Byun YS, Lee JH, Arsenault BJ, et al. Relationship of oxidized phospholipids on apolipoprotein B-100 to cardiovascular outcomes in patients treated with intensive versus moderate atorvastatin therapy: the TNT trial. *J Am Coll Cardiol* 2015;65:1286-95.
20. Byun YS, Yang X, Bao W, et al. Oxidized phospholipids on apolipoprotein B-100 and recurrent ischemic events following stroke or transient ischemic attack. *J Am Coll Cardiol* 2017;69:147-58.
21. Yeang C, Witztum JL, Tsimikas S. 'LDL-C'= LDL-C + Lp(a)-C: implications of achieved ultra-low LDL-C levels in the proprotein convertase subtilisin/kexin type 9 era of potent LDL-C lowering. *Curr Opin Lipidol* 2015;26:169-78.
22. Giugliano RP, Pedersen TR, Park JG, et al. Clinical efficacy and safety of achieving very low LDL-cholesterol concentrations with the PCSK9 inhibitor evolocumab: a prespecified secondary analysis of the FOURIER trial. *Lancet* 2017;390:1962-71.
23. Tsimikas S, Gordts PLSM, Nora C, Yeang C, Witztum JL. Statin therapy increases lipoprotein(a) levels. *Eur Heart J* 2019 May 20 (Epub ahead of print).
24. O'Donoghue ML, Fazio S, Giugliano RP, et al. Lipoprotein(a), PCSK9 inhibition, and cardiovascular risk. *Circulation* 2019;139:1483-92.
25. Leibundgut G, Scipione C, Yin H, et al. Determinants of binding of oxidized phospholipids on apolipoprotein (a) and lipoprotein (a). *J Lipid Res* 2013;54:2815-30.
26. van der Valk FM, Bekkering S, Kroon J, et al. Oxidized phospholipids on lipoprotein(a) elicit arterial wall inflammation and an inflammatory monocyte response in humans. *Circulation* 2016;134:611-24.
27. Scipione CA, Sayegh SE, Romagnuolo R, et al. Mechanistic insights into Lp(a)-induced IL-8 expression: a role for oxidized phospholipid modification of apo(a). *J Lipid Res* 2015;56:2273-85.
28. Que X, Hung MY, Yeang C, et al. Oxidized phospholipids are proinflammatory and proatherogenic in hypercholesterolaemic mice. *Nature* 2018;558:301-6.

Copyright © 2020 Massachusetts Medical Society.